

Spanish and Austrian goats were also in North America at that time. In 1904, the first dairy goat show was held in America at the World's Fair in St. Louis, Missouri. Also in 1904, the American Milk Goat Record Association was formed in Elyria, Ohio. Shortly thereafter, the association moved to North Carolina where it is currently located. In 1965, the name was changed to the American Dairy Goat Association (ADGA).

Today, goats provide a wide variety of products. They are important parts of the agriculture of many countries. Goat milk has a different fatty acid and protein structure than doe milk, and this often makes it easier for some people to digest. Individuals who are allergic to doe milk are commonly able to drink goat milk. Large parts of the world's population regularly drink goat versus doe milk. Worldwide, goat meat is consumed more than any other red meat and is a staple in many countries for festivals and holidays. Goats also provide two types of fiber—mohair and cashmere.

## Breeds of Goats

### Dairy Goats

The following descriptions have been adapted in part from the ADGA Breed Standards ([adga.org/breedstandards.html](http://adga.org/breedstandards.html)) and from the Dairy Herd Improvement Registry (DHIR) ([adga.org/DHIR/05breed\\_lactation\\_averages.htm](http://adga.org/DHIR/05breed_lactation_averages.htm)).

There are six breeds of dairy goats commonly found in the United States. These are the Alpine, LaMancha, Nubian, Oberhasli, Saanen, and Toggenburg. In 2005, two less common breeds, the Sable and the Nigerian Dwarf, were admitted into ADGA.

### Alpine



Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

The Alpine can be called either the French Alpine or the American Alpine, depending on parentage. However, when registered with ADGA, they are both referred to simply as Alpine. Alpines are a medium to large animal, have upright ears, medium to short hair, and a straight face. A Roman nose, Toggenburg color and markings, or an all white coat is discriminated against in the Alpine breed. Erect, medium-sized ears are preferred.

Alpines originated in the Alps region and are a fairly large and rangy goat. While no distinct color patterns have been bred for, there are typical color patterns that occur in the breed. Terms that describe the Alpine colors are as follows:

- Cou Blanc—Literally “white neck”; a white neck with black hindquarters and black or gray markings on the head.
- Cou Clair—Literally “clear neck”; a tan, saffron, off-white, or gray neck with black hindquarters.
- Cou Noir—Literally “black neck”; a black neck and front quarters with white hindquarters.



- Sundgau—Black with white markings on the underbody, facial stripes, and leg stripes, etc.
- Pied—Spotted or mottled in coloring.
- Chamoisee—Brown or bay with characteristic markings of a black face, dorsal stripe, feet and legs, and sometimes a martingale running over the withers and down the chest.
- Two-Tone Chamoisee—Light front quarters with brown or gray hindquarters. These animals are not Cou Blanc or Cou Clair as those animals have black hindquarters.
- Broken Chamoisee—A solid chamoisee color, broken by another color (usually white) by being banded or splashed.

Any variations in the above patterns broken with white should be described as a broken pattern, such as a broken Cou Blanc.

Alpines are known for being excellent milkers, with large, well-shaped udders. Alpines does should be at least 30 inches tall and weigh at least 135 pounds, while bucks must stand 32 inches tall and weigh a minimum of 170 pounds. According to ADGA and DHI, Alpines in 2005 averaged 2,334 pounds of milk, 3.3 percent milk fat, and 2.9 percent protein.

## LaMancha

The LaMancha is the only dairy goat that was developed in the United States. LaManchas are medium-sized animals, with any color combination acceptable. The hair is short and fine. The distinctive characteristic of the LaMancha is its very small ears. The history of the LaMancha is not well known. Short-eared goats are found throughout history and are thought to have originated in the United States from some goats that the Spanish missionaries brought to California.




Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

Two individuals seem to be responsible for developing the American LaMancha as it is seen today. The name LaMancha supposedly comes from a group of short-eared goats that were sent to the 1904 World's Fair in Paris for exhibition. The crate they were in was labeled "LaMancha, Cordoba, Spain." The name LaMancha was kept for these short-eared goats.

While Phoebe Wilhelm was supposedly the first to own a herd of 125 LaManchas in the early 1920s, Mrs. Eula Fay Frey from Oregon was very instrumental in developing today's LaMancha from some short-eared goats that were in a herd that she bought. She was amazed by how much milk these smaller animals produced. She bred these short-eared animals to Nubian and French Alpines and got short-eared offspring that were excellent milkers. She further developed the breed, and LaManchas were accepted as a breed into the ADGA registry on January 27, 1958, with Fay's Ernie, L-1. Approximately 200 animals were entered into the herd book at that time.


LaManchas have two types of ears—the "gopher ear" and the "elf ear." Gopher ears preferably have very little to no cartilage, with a maximum length of 1 inch (2.5 cm).



The end of the ear must be turned up or down. Bucks, or male goats, must have a gopher ear to be eligible for registration. The other type of ear is the “elf ear.” Elf ears are a maximum of 2 inches (5.0 cm) in length and the end of the ear must turn up or down. Bucks with elf ears are ineligible for registration.

Mature LaMancha does, or females, should stand 28 inches tall and weigh 130 pounds, while bucks should stand 30 inches tall and weigh a minimum of 160 pounds. According to ADGA and DHI, LaManchas in 2005 averaged 2,050 pounds of milk, 3.9 percent milk fat, and 3.1 percent protein.

## Nubian



Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

The Nubian is also called the Anglo-Nubian outside of the United States. It originated in England where bucks from the Nubia region of Africa and India were crossed with English does sometime around 1895. The Nubian is a large animal and is more heavily muscled than the other dairy breeds. This heavier muscling lends the Nubian breed to also being marketable as a meat goat.

Nubians are characterized by their long pendulous ears that must extend 1 inch

past the end of their muzzle, a Roman nose, and short, fine hair. They are well suited for hot weather conditions. Any color or color pattern is acceptable in the Nubian breed. Their milk is known for its higher milk fat and protein concentrations compared to the other dairy breeds; however, their milk production is usually lower.

Mature Nubian does should be at least 30 inches tall and weigh 135 pounds. Nubian bucks should be at least 32 inches tall and weigh at least 170 pounds. According to ADGA and DHI, Nubians in 2005 averaged 1,754 pounds of milk, 4.8 percent milk fat, and 3.7 percent protein.

## Oberhasli



Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

The Oberhasli breed is of Swiss origin and is a medium-sized animal. They are from the Brienzer region of Switzerland and are also known as the Oberhasli-Brienzer. When these animals were first imported to the United States in 1906 and again in 1920, they were called Swiss Alpines and were placed into the Alpine registry. In 1936, another group was imported to the United States. Today's Oberhaslis come from the 1936 imports. A small group of dedicated breeders



were determined to keep the Oberhasli as a pure breed and petitioned ADGA for its own herd book. In 1979, ADGA voted to make the Oberhasli its own breed, with its own herd book.

The Oberhasli differ from the Alpines in that they are slightly smaller in stature and more compact. They are characterized by their coloring, which is chamoisee. The females may be black, but the chamoisee coloring is preferred. The ADGA describes the standard coloring, called bay, as ranging from light to a deep red bay with the latter most desirable. A few white hairs through the coat and about the ears are permitted. Markings should include two black stripes down the face from above each eye to a black muzzle, a forehead that is nearly all black, black stripes from the base of each ear coming to a point just back of the poll and continuing along the neck and back as a dorsal stripe to the tail, a black belly and light tray to black udder, black legs below the knees and hocks, and ears that are black on the inside and bay on the outside. Bucks often have more black on the head than does, black whiskers, and black hair along the shoulder and lower chest with a mantle of black along the back. Bucks frequently have more white hairs throughout the coat than does. The face should be straight, as a Roman nose is discriminated against in the breed.

Mature height for an Oberhasli doe is 28 inches or more, and the mature weight should be near 120 pounds. Mature Oberhasli bucks must stand 30 inches tall and weigh 150 pounds. According to ADGA and DHI, the Oberhasli breed in 2005 averaged 2,137 pounds of milk, 3.4 percent milk fat, and 2.8 percent protein.


## Saanen




Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

Saanens originated in the Saanen Valley of Switzerland. Saanens are the largest of all the dairy breeds. Saanen coat color is either white or cream, with white being preferred. Spots on the skin are allowed and are actually common; however, spots on the hair coat are discouraged. The hair is short and fine and the face is straight or dished, with ears erect. Because of their light coloring, they are sensitive to sunlight and can sunburn easy. They tend to not perform as well in hot weather and prefer colder climates. Saanens have the highest milk production of all the dairy goat breeds, which has given them the nickname of “Queen of the Dairy Goats.”

Approximately 150 Saanens were imported between 1904 and the early 1930s from Switzerland. More Saanens came from England at a later date. Mature Saanen does should be at least 30 inches tall and weigh 135 pounds. The minimum height for mature bucks is 32 inches and 170 pounds. According to ADGA and DHI, the Saanen breed in 2005 averaged 2,537 pounds of milk, 3.3 percent milk fat, and 2.9 percent protein.



## Toggenburg




Courtesy of Curriculum Materials Service, The Ohio State University.  
Special thanks to Willow Run Dairy.

The Toggenburg originated in the Toggenburg Valley of Switzerland. It is the smallest and most compact of all the dairy breeds. Toggenburgs are a light fawn to a chocolate color with white markings. Their ears are white with a dark spot in the middle, and they have two white stripes down the face from above each eye to the muzzle. The hind legs are white from the hocks to the hooves, and the forelegs are white from the knees downward, with a dark vertical stripe below the knee being acceptable. They have a white triangle on either side of the tail, and white spots may be present at the root of the wattles or in the area where the wattles would be. The markings should be a light cream color, but white is the desired color. The ears are erect and carried straight forward, while the face is dished. As with the other breeds, except Nubian, a Roman nose is unacceptable. Toggenburgs have the longest hair of any of the dairy breeds, and it should be soft and very fine.

The Toggenburg is the smallest of the dairy breeds, with mature height for does being at least 26 inches. They should weigh 120 pounds. Mature bucks should be at least 28 inches tall with a minimum weight of 150

pounds. According to 2005 ADGA and DHI records, the Toggenburg breed averaged 2,101 pounds of milk, 3.3 percent milk fat, and 2.8 percent milk protein.

## Sable



Courtesy of International Sable Breeders Association.

The Sable is an animal of Saanen breeding that is not white. In Saanens, the gene for the white coat is dominant and a colored coat is recessive. When an animal of Saanen breeding is colored, it is because it has two recessive genes paired for hair coat color. When two Saanens with the recessive gene for hair color are mated, 25 percent of the offspring are likely to be colored. Colored Saanens are NOT eligible for registry in the Saanen herd book; however, they are eligible for registering in the Sable herd book. To register an animal in the Sable herd book, proof of Saanen, Sable, or Sable/Saanen parentage must be provided. Any color pattern for Sables is acceptable except solid white and solid light cream, and Toggenburg color is discouraged. Mature does must be at least 30 inches tall at the withers and weigh 135 pounds. Mature bucks must be at least 32 inches tall at the withers and weigh 170 pounds. The hair is short. The ears are erect and alertly carried, pointing forward. The face should be straight or dished.



## Nigerian Dwarf



Courtesy of Prairie Wood Ranch.

The Nigerian Dwarf is a small goat breed from Western Africa that has been developed for milk production more than for meat production. This goat is much smaller than the standard dairy goat breeds, standing 17 to 20 inches at the withers. Originally kept by African villagers to supply milk, these goats produce approximately 1 quart of high fat milk per day for up to 10 months.

The Nigerian Dwarf was imported to the United States in the 1980s, primarily to be exhibited at zoos and to be kept as pets. Their popularity grew due to their small size, ease of handling, and level of milk production.

The Nigerian Dwarf breeds at any time of the year and, therefore, can produce more than one kid crop per year. The doe should weigh 30 to 50 pounds, with a maximum height of 22.5 inches. The mature weight for a buck is 35 to 60 pounds, with a maximum height of 23.5 inches. Animals that are less than 17 inches tall at a mature age are discriminated against. The main colors of Nigerian Dwarfs are black, brown, or gold, with white markings. The Pygmy coloring, known as agouti, is discriminated against.

The Nigerian Dwarf produces approximately 2 to 4 pounds of milk per day, with anywhere

from 6 to 10 percent fat and 5 percent protein. These milk fat and protein concentrations are much higher than for the other dairy breeds, but remember, the Nigerian Dwarf produces much less milk overall. According to ADGA and DHI, the 2005 lactation average for the Nigerian Dwarf breed was 881 pounds of milk, with 6.5 percent fat and 4.0 percent protein.

## Meat Goats


Any breed of goat can be used for meat, and every goat should be looked at as a potential meat source. However, there are breeds that are more suited for meat production than others. This is usually because these breeds have been selected to produce bigger, heavier muscled, and leaner carcasses than other breeds. Several breeds commonly used for meat in the United States include Boer, Kiko, and Spanish goats. Often, the larger dairy breeds, such as Nubian, Alpine, and Saanen, are crossed with meat goats.

## Boer



Courtesy of Dail Gracy.

The Boer goat originated in South Africa and is thought to have originated from native African goats crossed with European goats that were brought by Dutch immigrants. The name "Boer" is a Dutch word for "farmer."



This term was used by the early Dutch settlers in the late 1800s to differentiate native goats from the Angora goats that were imported around the same time. In the early 1900s, South African farmers started to select animals for meat purposes and developed today's Boer goat. In 1959, a registry for Boer goats was established in South Africa. Boer goats were first imported into the United States in 1993 from Australia and New Zealand. The American Boer Goat Association was also formed in 1993 to record and certify these new Boer imports and Boer goat crosses.

Breed standards for the Boer goat include having a strong head with brown eyes, a Roman nose, and horns that are smooth and curve around the head (although animals that have been dehorned are not discriminated against). The ears are pendulous, similar to Nubians, and must hang smoothly against the side of the face. Boer goats traditionally have a white body with a red-brown head, or a red head with a white blaze on the face. Solid colors, other than all white or all black, are not discriminated against.

Animals should give the appearance of a meat animal with a broad chest, and a strong back, rump, and thighs. Mature Boer does weigh between 200 to 225 pounds, while mature Boer bucks weigh between 240 to 300 pounds. Kids are capable of gaining an average of 0.4 to 0.6 pounds per day. It is common to have a kidding rate of 200 percent in the herd. The Boer breed is known for extended breeding seasons, and it is possible to have three kiddings in two years.

Many commercial meat goat producers have found that Boer crosses perform exceptionally well or even better than purebred Boer goats, due to hybrid vigor. Crosses with the larger framed dairy goats, especially Nubians, are common.

## Kiko



Courtesy of Shelby Acres Okie-Dokie Kikos.

The Kiko goat was developed in New Zealand by a group of individuals who were interested in a meat animal that would perform well under natural browse conditions. Using native, feral goats and selecting specifically for production, the Kiko was developed. The word "kiko" is used by New Zealand native people for any meat-producing animal.

Specific traits were selected for during the development of this breed. Those traits included rapid growth rate, increased fertility (multiple births), early breeding age, good mothering skills with the ability to give birth unassisted, resistance to parasite infection, and soundness in feet. Animals were culled if they did not produce more than one kid, if they needed assistance with birthing, and if they showed any signs of developing foot rot or foot scald. The resulting breed is very hardy and adaptable to range living. Kids are noted for being extremely vigorous at birth and not needing any supplementation during growth.

Kikos are large animals. They are usually white, although they can be any color and any coat type (short or long). Kikos are excellent browsers and can survive quite well in open



rangeland under a wide variety of conditions and still have significant weight gains. Breed standards for Kikos include hardiness, capability of browsing for food, and maintaining substantial growth. Does should be able to conceive easily, carry multiple kids during pregnancy, and give birth without assistance. Mature males have a spiral horn, while females have smaller horns. Ears are high on the animal's head, are of moderate length, and are not pendulous or prick. The registered Kiko is typically identified with a microchip or tattoo, plus an ear tag.

### Spanish Goat



Photo by Phillip Sponenberg.  
Courtesy of the American Livestock Breeds Conservancy.

The current-day Spanish goat is thought to be a descendant of the goats Spanish explorers brought to the United States as meat and milk producers. They are not of any specific breed but describe a type of goat. Spanish goats are smaller, meaty animals that browse well on range conditions. They typically have small udders so that they do not get scratched up by brush and cacti. They can be any color, horned or not horned, and have different types of hair coats. Some Spanish goats are also used for cashmere. Most of the Spanish goats in the United States are located in Texas, are raised on the range, and are selected for meat production.

## Fiber Goats

### Angora



Courtesy of Lavender Hill Farm.

Angora goats are raised primarily for the mohair, or fiber, from their fleece. Angora goats are excellent producers of fine quality fleece. If cared for properly, an Angora goat can produce anywhere from 8 to 16 pounds per year of mohair. They are thought to have originated in the Himalayan Mountains of Asia. Angora goats migrated to Turkey sometime during the 13th century. Their name comes from the Ankara (Angora, prior to 1930) province in Turkey, where they were first bred. Although these goats were not exported from Turkey to the United States until 1849, the United States is one of the principal mohair producers in the world today, producing more than 13 million pounds of mohair each year. Turkey and South Africa are two other major mohair producers in the world. Angora goats are known for being docile, compared with other breeds of goats, but also for being "delicate." The fleece grows year-round at the rate of  $\frac{3}{4}$  to 1 inch per month. This fast rate of growth puts a large amount of strain on the animal and may contribute to their relative lack of hardiness. At mature weight, does weigh





approximately 75 to 100 pounds and wethers (castrated males) and bucks weigh between 85 to 125 pounds.

## Cashmere



Courtesy of Liberty Farm Cashmere.

Cashmere goats are a type, not a breed, so there is no such thing as a “purebred cashmere” goat. Cashmere is the fine underdown from goats, which most goats (except Angora) produce in varying quantities. A “cashmere goat” is one that produces commercial quantities of cashmere. The fleece from a cashmere goat consists of very fine, crimped down and longer, coarse outside guard hairs. The down undercoat is grown during late summer and autumn and is shed naturally in the spring. Cashmere fibers can be separated from the fleece either by combing out the down from the animal or by means of a commercial dehairer on sheared fleeces. In the United States, Spanish meat goats from Texas and the southwest are used for cashmere breeding stock, in addition to being used for meat production. Of the dairy breeds, Toggenburg, Saanen, and Nubian also produce cashmere.

## Other Breeds

### Pygmy Goat



Department of Animal Sciences, The Ohio State University.


The Pygmy goat (originally called the Cameroon Dwarf Goat) was originally developed in the French Cameroon area of western Africa. This goat is very small, with a full-grown Pygmy goat standing 16 to 23 inches at the withers. Females are slightly smaller than males. The most common coat color is known as “agouti.” Agouti means that the hairs are two-toned, giving a salt and pepper appearance to the coat. Pygmy goats are very hardy and gregarious. They also have a good temperament and are excellent with children.

The first U.S. Pygmy goats were imported from Sweden in 1959. Since that time, they have grown in popularity for use in 4-H and FFA projects and in children’s zoos. Some of their growth in popularity has occurred because they require less space and feed.


Pygmy goats have also been used for family milk and meat production. A doe in milk can be expected to give about 1 pint of high-fat milk twice a day for 4 to 6 months. The carcass is well muscled, providing meat similar in quantity to a spring lamb. The Pygmy goat, unlike most goat breeds, breeds




# Goat Hoof Trimming




## Tools



Foot Rot Shears




Pruning Shears




Overgrown Hoof



Cut from Heel to Toe



Properly Trimmed Hoof Bottom View




Properly Trimmed Hoof



**Exploratory Learning: Educational Program**  
This component adapted from materials of the American Dairy Goat Association

Product distribution through the Curriculum Materials Service





### Intramuscular injection sites

Yes, correct injection site.

No, don't inject here.

Intramuscular injections are commonly given in the triangular area of the neck, bounded caudally by the shoulder, dorsally by the cervical vertebra, and ventrally by the nuchal ligament of the spinous process. Do not give intramuscular injections on the round for goats. Volume given in the muscle should not be more than 3 milliliters per site.

Ohio State University Extension.

- ~~4. Give injections at clean, dry sites on the animal.~~
- ~~5. Do not transfer needles back and forth from animal to bottle. You may carry bacteria from the animal's skin. Needles are used just once and then discarded; do not use the same syringe needle multiple times.~~
- ~~6. Products with low dosage rates are preferred.~~
- ~~7. When making multiple injections, space injections at least six inches apart.~~

### Subcutaneous injection site

Subcutaneous injections are usually given in the auxiliary region behind the elbow joint or in the triangular area of the neck in front of the shoulders.

Ohio State University Extension.

#### **Injection Reminders**

All products labeled SQ are administered behind the elbow joint or in front of the shoulder. Without exception and regardless of age of the animal, all products labeled IM are administered in the neck region.

Whenever possible, IM injections are to be avoided. Products labeled for SQ, IV, or O administration are recommended.



*No figuring withdrawal time*

Use this poster in conjunction with Medication Label - Read the Following Drug Label and/or Insert, and Correctly Match Each Part to Its Corresponding Place situation/task statement and identification tags.

# Medication Label

Name of Drug \_\_\_\_\_

**OMNIBIOTIC**

(hydrocillin)

Active Ingredients \_\_\_\_\_

**Directions for use: See package insert**

Cautions and Warnings \_\_\_\_\_

Withholding Times \_\_\_\_\_

**Warning:** The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food. Exceeding the highest recommended dosage level may result in antibiotic residues in meat or milk beyond the withdrawal time.

Quantity of Contents \_\_\_\_\_

Net Contents: 100 ml

Distributed by

Name of Distributor \_\_\_\_\_

**USA Animal Health, Inc.**

Store between 2° and 8° C (36° and 46° F). \_\_\_\_\_ Storage

**Keep dry and away from light.**

TAKE TIME



OBSERVE LABEL DIRECTIONS



Exploratory Learning: Educational Program

Product distributed through the Curriculum Materials Service