


Use this poster in conjunction with Anatomy - Match the correct part names with their locations on the diagram situation/task statement and/or Parts of a Horse Identification tags.

Parts of a Horse



Exploratory Learning: Educational Program

Product distribution through Curriculum Materials Service

Example

Blue Bird

Horse 12
For mature horses

Guaranteed analysis

Crude protein min12%
Crude fat min 3.0%
Crude fiber max10.0%
Calcium min .8% max 1.20%
Phosphorus min60%
Salt min .8% max 1.10%
Copper min 60 ppm
Zinc min 300 ppm
Selenium min3 PPM
Vitamin A min 10,000 IU/lb

Ingredients

Rollod Corn, Rolled Oats, Rolled Barley, wheat mill run, Soybean Meal, Salt, Cane Molasses, Calcium Carbonate, Vitamin A supplement, Vitamin D3 Supplement, Vitamin E Supplement, Manganese Oxide, Zinc Oxide, Copper Sulfate, Magnesium Oxide, Cobalt Carbonate, Sodium Selenate.

Feeding directions

Feed up to 1% of body weight per head per day along with at least 1% of body weight of hay or pasture. Provide salt and plenty of fresh, clean water at all times. Make all feed changes gradually. This feed is balanced to be fed with local grass hays. Feed ALFA-HORSE 8 if feeding any alfalfa hay.

Do not feed to foals under 12 months old. Do not feed moldy or infested feeds.

Manufactured by:

Blue Bird Feeds
16 Main St Bluebird, OR 97999 Net wt 50 lb

1. Who manufactured this Feed?
Blue Bird Feeds
2. What is the minimum amount of copper available in this feed?
60 ppm
3. What age of horses is this feed fed to?
Mature horses
4. What is the main ingredient in this feed?
Rollod Corn
5. What age of horses should not get this feed?
Foals under 12 months
6. Is Soybean Meal in this feed?
Yes

Use this poster in conjunction with Care - Read the following drug label and/or insert, and correctly match each part to its corresponding location *situation/task statement* and/or Medication Label/Insert *identification tags*.

Medication Label

Name of Drug _____ **Active Ingredients** _____

Equibiotic
(hydrocillin)

Directions for use: See package insert

Warning: The use of this drug must be discontinued for 30 days before treated animals are slaughtered for food. Exceeding the highest recommended dosage level may result in antibiotic residues in meat or milk beyond the withdrawal time.

Withholding Times _____

Storage _____

Store between 2° and 8° C (36° and 46° F)
Keep dry and keep away from light

Quantity of Contents _____ **Net contents: 100 ml**
TAKE TIME
Distributed by
USA Animal Health, Inc.

Name of Distributor _____

Cautions and Warnings _____

Quantity of Contents _____

TAKE TIME

OBSERVE LABEL DIRECTIONS



Exploratory Learning: Educational Program

This component relates to information found in Unit 5, Level 1 of *Equine Awareness Youth Education Program: Educators' Curriculum Guide*, (Green, Schmitt, and Whiteaker, 1997).

Product distribution through Curriculum Materials Service




Figure 104. The neck is the safest injection site. Give the injection within the triangular area. If multiple injections are needed, use the other sites shown in the following figures in order to decrease soreness on the neck.



Figure 105. Butt.



Figure 106. Chest.